

**Clinical description, Revised 2004.** [http://www.cdc.gov/epo/dphsi/casedef/arboviral\\_current.htm](http://www.cdc.gov/epo/dphsi/casedef/arboviral_current.htm)

### **West Nile virus disease, and other domestic neurotropic arboviral diseases**

Arboviral infections may be asymptomatic or may result in febrile illnesses of variable severity sometimes associated with central nervous system (CNS) involvement. When the CNS is affected, clinical syndromes include aseptic meningitis, myelitis and encephalitis, which are clinically indistinguishable from similar syndromes caused by other viruses. Arboviral meningitis is usually characterized by fever, headache, stiff neck, and pleocytosis in cerebrospinal fluid. Arboviral myelitis is usually characterized by fever and acute bulbar or limb paresis or flaccid paralysis. Arboviral encephalitis is usually characterized by fever, headache, and altered mental status ranging from confusion to coma with or without additional signs of brain dysfunction. Less common neurological syndromes can include cranial and peripheral neuritis or other neuropathies, including Guillain-Barré syndrome.

Non-neuroinvasive syndromes caused by these usually neurotropic arboviruses can rarely include myocarditis, pancreatitis, or hepatitis. In addition, they may cause febrile illnesses (e.g., West Nile fever [WNF]) that are non-localized, self-limited illnesses with headache, myalgias, arthralgias, and sometimes accompanied by skin rash or lymphadenopathy. Laboratory-confirmed arboviral illnesses lacking documented fever can occur, and overlap among the various clinical syndromes is common.

### **Clinical criteria for diagnosis**

Cases of arboviral disease are classified either as neuroinvasive or non-neuroinvasive, according to the following criteria:

Neuroinvasive disease requires the presence of fever and at least one of the following, as documented by a physician and in the absence of a more likely clinical explanation:

- Acutely altered mental status (e.g., disorientation, obtundation, stupor, or coma), or
- Other acute signs of central or peripheral neurologic dysfunction (e.g., paresis or paralysis, nerve palsies, sensory deficits, abnormal reflexes, generalized convulsions, or abnormal movements), or
- Pleocytosis (increased white blood cell concentration in cerebrospinal fluid [CSF]) associated with illness clinically compatible with meningitis (e.g., headache or stiff neck).

Non-neuroinvasive disease requires, at minimum, the presence of documented fever, as measured by the patient or clinician, the absence of neuroinvasive disease (above), and the absence of a more likely clinical explanation for the illness. Involvement of non-neurological organs (e.g., heart, pancreas, liver) should be documented using standard clinical and laboratory criteria.

### **Laboratory criteria for diagnosis**

Cases of arboviral disease are also classified either as confirmed or probable, according to the following laboratory criteria:

Confirmed case :

- Four-fold or greater change in virus-specific serum antibody titer, or
- Isolation of virus from or demonstration of specific viral antigen or genomic sequences in tissue, blood, CSF, or other body fluid, or
- Virus-specific immunoglobulin M (IgM) antibodies demonstrated in CSF by antibody-capture enzyme immunoassay (EIA), or
- Virus-specific IgM antibodies demonstrated in serum by antibody-capture EIA and confirmed by demonstration of virus-specific serum immunoglobulin G (IgG) antibodies in the same or a later specimen by another serologic assay (e.g., neutralization or hemagglutination inhibition).

Probable case :

- Stable (less than or equal to a two-fold change) but elevated titer of virus-specific serum antibodies, or
- Virus-specific serum IgM antibodies detected by antibody-capture EIA but with no available results of a confirmatory test for virus-specific serum IgG antibodies in the same or a later specimen.

**Case definition**

A case must meet one or more of the above clinical criteria and one or more of the above laboratory criteria.

**Comment**

Because closely related arboviruses exhibit serologic cross-reactivity, positive results of serologic tests using antigens from a single arbovirus can be misleading. In some circumstances (e.g., in areas where two or more closely related arboviruses occur, or in imported arboviral disease cases), it may be epidemiologically important to attempt to pinpoint the infecting virus by conducting cross-neutralization tests using an appropriate battery of closely related viruses. This is essential, for example, in determining that antibodies detected against St. Louis encephalitis virus are not the result of an infection with West Nile (or dengue) virus, or vice versa, in areas where both of these viruses occur. Because dengue fever and West Nile fever can be clinically indistinguishable, the importance of a recent travel history and appropriate serologic testing cannot be overemphasized. In some persons, West Nile virus-specific serum IgM antibody can wane slowly and be detectable for more than one year following infection. Therefore, in areas where West Nile virus has circulated in the recent past, the co-existence of West Nile virus-specific IgM antibody and illness in a given case may be coincidental and unrelated. In those areas, the testing of serially collected serum specimens assumes added importance.

The seasonality of arboviral transmission is variable and depends on the geographic location of exposure, the specific cycles of viral transmission, and local climatic conditions. Reporting should be etiology-specific (see below; the six diseases printed in bold are nationally reportable to CDC):

- **St. Louis encephalitis virus disease**
- **West Nile virus disease**
- **Powassan virus disease**
- **Eastern equine encephalitis virus disease**
- **Western equine encephalitis virus disease**
- **California serogroup virus disease** (includes infections with the following viruses: California encephalitis, Jamestown Canyon, Keystone, La Crosse, snowshoe hare, and trivittatus)